arXiv:1407.0839v2 [math.PR] 20 Oct 2014

AN EXPONENTIAL INEQUALITY FOR SYMMETRIC RANDOM VARIABLES

Raphaël Cerf
ENS Paris

and

Matthias Gorny
Université Paris Sud and ENS Paris

A central question in the study of sums of independent identically distributed random variables is to control the probabilities that they deviate from their typical values. We present here a very simple exponential inequality. This inequality holds for any symmetric distribution, and does not require any integrability condition.

Theorem. *Let $n \geq 1$ and let $X_1, \ldots, X_n$ be n independent identically distributed symmetric real-valued random variables. For any $x, y > 0$, we have*

$$\mathbb{P}\big(X_1 + \cdots + X_n \geq x,\, X_1^2 + \cdots + X_n^2 \leq y\big) < \exp\left(-\frac{x^2}{2y}\right).$$

If we apply this inequality with nx, ny instead of x, y, we obtain an inequality controlling the first two empirical moments of the sequence $X_1, \cdots, X_n$:

$$\mathbb{P}\left(\frac{X_1 + \cdots + X_n}{n} \geq x,\, \frac{X_1^2 + \cdots + X_n^2}{n} \leq y\right) < \exp\left(-\frac{nx^2}{2y}\right).$$

With the classical theory of large deviations [3], we usually obtain exponential inequalities of this type, but unfortunately they are valid for n large and it is a difficult task to quantify how large n has to be. During the last decades, probabilists and statisticians have been trying to find non asymptotic inequalities, which are valid for any $n \geq 1$ and with explicit constants [1]. Most of these inequalities are based on the phenomenon of concentration of measure and they require a strong control on the tail of the distribution of X_1, typically the existence of an exponential moment. The inequality we present here is a deviation inequality which is valid for all $n \geq 1$. We suppose that the distribution is symmetric, and no further integrability condition is required. Initially, this inequality was obtained through classical results of large deviations within Cramér's theory and an inequality on the rate function derived to study a Curie–Weiss model of self–organized criticality [2]. Since the statement of the exponential inequality is very simple, we looked for an elementary proof, and it is this proof we present here.

Let us prove the theorem. We suppose that $\mathbb{P}(X_1 = 0) < 1$, otherwise the inequality of the theorem is immediate. Let $n \geq 1$ and $x, y > 0$. We set

$$S_n = X_1 + \cdots + X_n\,, \qquad T_n = X_1^2 + \cdots + X_n^2\,.$$

Let $s, t > 0$. We have

$$\begin{aligned}\mathbb{P}\big(S_n \geq x,\, T_n \leq y\big) &= \mathbb{P}\left(sS_n \geq sx,\, -tT_n \geq -ty\right)\\ &\leq \mathbb{P}\big(sS_n - tT_n \geq sx - ty\big)\\ &\leq \mathbb{P}\Big(\exp\big(sS_n - tT_n\big) \geq \exp\big(sx - ty\big)\Big)\,.\end{aligned}$$

We recall one of the most classical stochastic inequality.

Markov's inequality. If X is a non–negative random variable, then

$$\forall \lambda > 0 \qquad \mathbb{P}(X \geq \lambda) \leq \frac{\mathbb{E}(X)}{\lambda}\,.$$

Using Markov's inequality and the fact that $X_1, \ldots, X_n$ are i.i.d., we get

$$\begin{aligned}\mathbb{P}\big(S_n \geq x,\, T_n \leq y\big) \,&\leq\, \exp\big(-sx + ty\big)\,\mathbb{E}\left(\prod_{i=1}^{n}\exp\big(sX_i - tX_i^2\big)\right)\\ &=\, \exp\big(-sx + ty\big)\left(\mathbb{E}\Big(\exp\big(sX_1 - tX_1^2\big)\Big)\right)^n\,.\end{aligned}$$

The distribution of X_1 is symmetric thus

$$\begin{aligned}\mathbb{E}\Big(\exp\big(sX_1 - tX_1^2\big)\Big) \,&=\, \mathbb{E}\Big(\exp\big(-sX_1 - tX_1^2\big)\Big)\\ &=\, \frac{1}{2}\Big(\mathbb{E}\Big(\exp\big(sX_1 - tX_1^2\big)\Big) + \mathbb{E}\Big(\exp\big(-sX_1 - tX_1^2\big)\Big)\Big)\\ &=\, \mathbb{E}\Big(\cosh(sX_1)\,\exp\big(-tX_1^2\big)\Big)\,.\end{aligned}$$

We choose now $t = s^2/2$. We have the inequality

$$\forall u \in \mathbb{R}\backslash\{0\} \qquad \cosh(u)\,\exp\big(-u^2/2\big) < 1\,.$$

Since $\mathbb{P}(X_1 = 0) < 1$, the above inequality implies that

$$\mathbb{E}\Big(\cosh(sX_1)\,\exp\big(-s^2X_1^2/2\big)\Big) \,<\, 1\,,$$

whence also

$$\mathbb{P}\big(S_n \geq x,\, T_n \leq y\big) \,<\, \exp\big(-sx + s^2y/2\big)\,.$$

We finally choose $s = x/y$ and we obtain the desired inequality.

References

[1] Stéphane Boucheron, Gàbor Lugosi, and Pascal Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 2013.

[2] Raphaël Cerf and Matthias Gorny. A Curie-Weiss model of Self-Organized Criticality. *The Annals of Probability*, to appear, 2013.

[3] Amir Dembo and Ofer Zeitouni. *Large deviations techniques and applications*, volume 38 of *Stochastic Modelling and Applied Probability*. Springer-Verlag, 2010.